

doi: 10.13241/j.cnki.pmb.2020.11.019

CRP、PCT、WBC 联合检测对中老年社区获得性肺炎的诊断效果*

刘欢^{1,2} 朱述阳^{3Δ} 于海容² 洪倩⁴ 潘倩⁵

(1 徐州医科大学 江苏 徐州 221004; 2 徐州医科大学附属淮海医院呼吸科 江苏 徐州 221000;

3 徐州医科大学附属医院呼吸科 江苏 徐州 221000; 4 徐州医科大学附属淮海医院儿科 江苏 徐州 221000;

5 徐州医科大学附属淮海医院检验科 江苏 徐州 221000)

摘要 目的:研究 C 反应蛋白(C-reactive protein, CRP)、降钙素原(Procalcitonin, PCT)以及白细胞(White blood cell, WBC)联合检测对中老年社区获得性肺炎患者的诊断效果。**方法:**选择 2018 年 1 月~2019 年 1 月我院收治的 76 例中老年社区获得性肺炎患者为观察组,同期在我院体检中心选取 30 例健康体检者为对照组。检测两组研究对象的 CRP、PCT 及 WBC 水平。采取肺炎严重程度 CURB-65 评分将观察组的 76 例患者分为低危组($n=63$ 例), CURB-65 评分 <3 分,以及高危组($n=13$ 例), CURB-65 评分 ≥ 3 分;依照观察组的转归情况分为存活组($n=70$ 例)以及死亡组($n=6$ 例),比较观察组和对照组患者,低危组和高危组患者,存活组和死亡组患者的 CRP、PCT 及 WBC 水平的差异。**结果:**观察组患者的 CRP、PCT 及 WBC 水平明显高于对照组($P<0.05$);高危组患者的 CRP、PCT 水平明显高于低危组患者($P<0.05$),而 WBC 水平两组无明显差异($P>0.05$);死亡组患者的 CRP、PCT 水平明显高于存活组患者($P<0.05$),而 WBC 水平两组无明显差异($P>0.05$)。经 Pearson 相关分析发现, CURB-65 评分与 PCT、CRP 均呈明显的正相关($t=0.532, 0.497, P$ 均 <0.05)。**结论:**CRP、PCT 以及联合检测可以为中老年社区获得性肺炎的诊断提供有利的信息,且 CRP、PCT 与患者的病情严重程度有一定的相关性。

关键词:C 反应蛋白;降钙素原;白细胞;中老年社区获得性肺炎

中图分类号:R563.1 **文献标识码:**A **文章编号:**1673-6273(2020)11-2094-04

Diagnostic Efficacy of Combined Detection of CRP, PCT and WBC in Elderly Patients with Community Acquired Pneumonia*

LIU Huan^{1,2}, ZHU Shu-yang^{3Δ}, YU Hai-rong², HONG Qian⁴, PAN Qian⁵

(1 Xuzhou Medical University, Xuzhou, Jiangsu, 221004, China; 2 Department of Respiratory, Huaihai Hospital, Xuzhou Medical University, Xuzhou, Jiangsu, 221000, China; 3 Department of Respiratory, Affiliated Hospital of Xuzhou Medical University, Xuzhou, Jiangsu, 221000, China; 4 Department of Pediatrics, Huaihai Hospital, Xuzhou Medical University, Xuzhou, Jiangsu, 221000, China;

5 Department of Clinical Laboratory, Huaihai Hospital, Xuzhou Medical University, Xuzhou, Jiangsu, 221000, China)

ABSTRACT Objective: To investigate the diagnostic efficacy of combined detection of C-reactive protein (CRP), procalcitonin (PCT) and white blood cell (WBC) in elderly patients with community acquired pneumonia. **Methods:** Selected 76 cases of elderly patients with community acquired pneumonia who were treated in our hospital from January 2018 to January 2019 as observation group. In the same period, 30 healthy people in our physical examination center were selected as the control group. The levels of CRP, PCT and WBC in the two groups were detected. The CURB-65 score of pneumonia was used to classify 76 patients in the observation group into low-risk group ($n=63$ cases), CURB-65 score <3 points, and high-risk group ($n=13$ cases), CURB-65 score ≥ 3 points; according to the outcome of the observation group, the survival group ($n=70$ cases) and the death group ($n=6$ cases), the comparison observation group and the control group, the low-risk group and the high-risk group, the survival group and the death group. Differences in CRP, PCT and WBC levels in the group. **Results:** The CRP, PCT and WBC levels in the observation group were significantly higher than control group ($P<0.05$). The levels of CRP and PCT in the high-risk group were significantly higher than those in the low-risk group ($P<0.05$), but there was no significant difference between the two groups ($P>0.05$). The CRP and PCT levels in the death group were significantly higher than those in the surviving group($P<0.05$), but there was no significant difference between the two groups ($P>0.05$). Pearson correlation analysis showed that CURB-65 score was positively correlated with PCT and CRP ($t=0.532, 0.497, P<0.05$). **Conclusion:** The combined detection of CRP and PCT can provide useful information for the diagnosis of community-acquired pneumonia in the elderly, and has a certain correlation with the severity of the disease.

Key words: C-reactive protein; Procalcitonin; White blood cell; Elderly community-acquired pneumonia

Chinese Library Classification(CLC): R563.1 **Document code:** A

Article ID: 1673-6273(2020)11-2094-04

* 基金项目:国家自然科学基金青年基金项目(81500951)

作者简介:刘欢(1988-),女,硕士研究生,主治医师,研究方向:呼吸方向,电话:15862149873, E-mail:lhshitou2019@163.com

Δ 通讯作者:朱述阳(1963-),男,硕士,主任医师,研究方向:呼吸方向,电话:18052268665, E-mail:jiechenz356@sohu.com

(收稿日期:2019-12-01 接受日期:2019-12-23)

前言

社区获得性肺炎指的是在医院外患有的一种感染性肺部实质炎症疾病,并且包括入院之后在平均潜伏期间发病的情况,主要是由于非典型病原体、细菌或者病毒等感染而引发^[1-3]。近年来,由于抗菌药物的滥用和病原体的不断变迁,我国社区获得性肺炎的发病率和病死率迅速升高^[4,5]。研究发现中老年人是社区获得性肺炎发病和病死的高危人群^[6-8]。早期发现、及时确诊、有效治疗和准确评估疾病严重程度,对于改善中老年社区获得性肺炎的预后极为关键。寻找快速、特效、敏感和简便的早期诊断和评估病情的生化检测指标,是目前诊疗中老年社区获得性肺炎亟待解决的重要问题^[9,10]。中老年社区获得性肺炎患者出现咳嗽、咳痰以及发热等典型的肺炎症状通常不明显,而主要表现为乏力、纳差以及晕厥等情况,导致临床医师对病情的评估具有较大的局限性^[11]。社区获得性肺炎患者的肺实质会发生炎症性病变,准确鉴别病因是开展有效治疗的关键,CRP、PCT 以及 WBC 均属于机体的炎症标志物。本研究采用 CRP、PCT 以及 WBC 联合检测,以分析其对中老年社区获得性肺炎的诊断效果。

1 资料与方法

1.1 一般资料

选择 2018 年 1 月~2019 年 1 月我院收治的 76 例中老年社区获得性肺炎患者为观察组,同期在我院体检中心选取 30 例健康体检者为对照组。纳入标准:(1)年龄大于 18 岁;(2)符合中华医学会呼吸病学会 2016 年制定的《社区获得性肺炎诊断和治疗指南》的诊断标准。排除标准:(1)免疫缺陷疾病;

(2)长期使用免疫抑制剂或激素治疗;(3)怀疑活动性肺结核、肺部肿瘤患者。观察组 76 例,男 41 例,女 35 例;年龄 45~73 岁,平均(63.38±6.54)岁。对照组 30 例,男 16 例,女 14 例;年龄 45~72 岁,平均(61.26±5.39)岁。

1.2 方法

所有患者均抽取 5 mL 空腹静脉血,采取乳胶免疫比浊法检测血清 CRP 水平,试剂盒购自深圳迈瑞生物医疗电子股份有限公司;采取酶联免疫荧光法检测血清 PCT 水平,试剂盒购自生物梅里埃法国股份有限公司;采取全自动血液细胞分析仪(深圳迈瑞生物医疗电子股份有限公司)检测 WBC 水平。

采取肺炎严重程度 CURB-65 评分将观察组的 76 例患者分为低危组(n=63 例),CURB-65 评分<3 分,以及高危组(n=13 例),CURB-65 评分≥3 分;依照观察组的转归情况分为存活组(n=70 例)以及死亡组(n=6 例),比较观察组和对照组,低危组和高危组,存活组和死亡组 CRP、PCT 以及 WBC 水平的差异。并采取 Pearson 相关分析检测 CURB-65 评分与 CRP、PCT 以及 WBC 的相关性。

1.3 统计学分析

采用 SPSS 21.0,计量资料以 $\bar{x} \pm s$ 表示,组间和组内对比用方差分析和 t 检验,组间率的比较用 χ^2 检验, $P < 0.05$ 为差异有统计学意义。

2 结果

2.1 两组患者的 CRP、PCT 以及 WBC 水平对比

观察组患者的 CRP、PCT 以及 WBC 水平明显高于对照组患者($P < 0.05$),见表 1。

表 1 两组患者的 CRP、PCT 以及 WBC 水平对比($\bar{x} \pm s$)

Table 1 Comparison of CRP, PCT and WBC levels between the two groups($\bar{x} \pm s$)

Groups	n	CRP(mg/L)	PCT(ng/L)	WBC($\times 10^9/L$)
Control group	30	5.37±1.26	0.13±0.02	6.34±1.57
Observation group	76	59.48±11.39*	1.65±0.44*	13.29±2.78*

Note: Compared with the control group, * $P < 0.05$.

2.2 高危组和低危组患者的 CRP、PCT 以及 WBC 水平对比 ($P < 0.05$),而 WBC 水平两组无明显差异($P > 0.05$)见表 2。

高危组患者的 CRP 和 PCT 水平明显高于低危组患者

表 2 高危组和低危组患者的 CRP、PCT 以及 WBC 水平对比($\bar{x} \pm s$)

Table 2 Comparison of CRP, PCT and WBC levels in high-risk and low-risk groups($\bar{x} \pm s$)

Groups	n	CRP (mg/L)	PCT (ng/L)	WBC($\times 10^9/L$)
Low-risk group	63	30.42±5.17	1.14±0.25	13.63±3.57
High-risk group	13	79.36±12.49*	2.13±0.59*	14.38±3.49

Note: Compared with the control group, * $P < 0.05$.

2.3 存活组和死亡组患者的 CRP、PCT 以及 WBC 水平对比

死亡组患者的 CRP 和 PCT 水平明显高于存活组患者($P < 0.05$),而 WBC 水平两组无明显差异($P > 0.05$),见表 3。

2.4 相关性分析

经 Pearson 相关分析发现,CURB-65 评分与 PCT、CRP 均

呈明显的正相关($t=0.532, 0.497, P$ 均 < 0.05)。

3 讨论

目前,中老年社区获得性肺炎发病时的病情往往较为严重,病死率和并发症率比较高^[12-14]。虽然抗生素在临床上得到了

日益广泛的应用,社区获得性肺炎依旧是威胁人类生命健康的一种重要感染性疾病^[15]。在中老年人中具有极高的发病率,分析其原因,一方面,中老年人呼吸道的防御能力随着年龄的増长而明显下降,另一方面,中老年患者的呼吸道症状大多并不突出,加上患者大多合并有一种甚至多种的基础性疾病,胸片

检查可能由于其他疾病而不显著,极易耽误病情,而且一旦中老年人发生肺部感染,更易迅速发展成重症肺炎^[16-19]。以往临床诊断肺部感染主要参考患者的白细胞分类及计数、症状表现、痰细菌培养结果和胸部 X 线检查,但是上述指标的特异度及敏感度均较低^[20]。

表 3 存活组和死亡组患者的 CRP、PCT 以及 WBC 水平对比($\bar{x} \pm s$)

Table 3 Comparison of CRP, PCT and WBC levels in patients in the survival and death groups($\bar{x} \pm s$)

Groups	n	CRP (mg/L)	PCT (ng/L)	WBC($\times 10^9/L$)
Survival group	70	34.92 $\pm$ 5.63	1.35 $\pm$ 0.37	14.64 $\pm$ 3.57
Dead group	6	85.42 $\pm$ 13.79*	2.97 $\pm$ 0.84*	15.29 $\pm$ 3.64

Note: Compared with the control group, * $P < 0.05$.

CRP 作为一种典型的炎症标志物,其水平的升高表明机体出现了炎症反应。CRP 能对机体中的补体发挥激活功能,从而使吞噬细胞的作用増强,强化吞噬细胞在体内的清理作用,发挥清除病原微生物以及保护机体组织的目的^[21-23]。PCT 作为感染性疾病的一种血清学标志物,具有高特异性以及高灵敏性的优点。当出现严重的寄生虫、细菌、真菌感染,脓毒症和多脏器功能衰竭时,机体的血清 PCT 水平会大幅度升高,而且其水平不会受到内部激素水平的影响^[24]。有研究发现,PCT 在鉴别炎性反应和细菌感染方面较体温、WBC 和 CRP 具有更好的临床应用价值^[25]。血清 PCT 水平不仅可以反映患者全身炎症反应的活跃程度,还可以反映被感染器官的类型、大小、患者炎症的严重程度、感染细菌的种类以及患者自身免疫反应情况等^[26-30]。本研究结果发现,观察组患者的 CRP、PCT 以及 WBC 水平明显高于对照组;高危组患者的 CRP、PCT 水平明显高于低危组患者;死亡组患者的 CRP、PCT 水平明显高于存活组患者;表明 CRP、PCT 水平越高,中老年社区获得性肺炎患者的病情就越严重,预后就越不佳。本研究经 Pearson 相关分析发现,CURB-65 评分与 PCT、CRP 均呈明显的正相关。表明 CRP、PCT 水平越高,中老年社区获得性肺炎患者的 CURB-65 评分就越高。CRP、PCT 水平的高低对于评估中老年社区获得性肺炎病情严重程度和疾病转归具有重要的指导意义。

综上所述,CRP、PCT 联合检测可以为中老年社区获得性肺炎的诊断提供有利的信息,且与患者的病情严重程度有一定的相关性。

参考文献(References)

- [1] Curbelo J, Luquero BS, Jm NRN, et al. Inflammation biomarkers in blood as mortality predictors in community-acquired pneumonia admitted patients: Importance of comparison with neutrophil count percentage or neutrophil-lymphocyte ratio [J]. Plos One, 2017, 12(3): e0173947
- [2] Lópezdeandrés A, Migueldíez JD, Jiméneztrujillo I, et al. Hospitalisation with community-acquired pneumonia among patients with type 2 diabetes: an observational population-based study in Spain from 2004 to 2013[J]. Bmj Open, 2017, 7(1): e013097
- [3] Domínguez À, Soldevila N, Toledo D, et al. Effectiveness of 23-valent pneumococcal polysaccharide vaccination in preventing community-acquired pneumonia hospitalization and severe outcomes in the elderly in Spain[J]. Plos One, 2017, 12(2): e0171943
- [4] Saghaian-Hedengren S, Mathew JL, Hagel E, et al. Assessment of Cy-

- tokine and Chemokine Signatures as Potential Biomarkers of Childhood Community-acquired Pneumonia Severity: A Nested Cohort Study in India[J]. Pediatr Infect Dis J, 2017, 36(1): 102-108
- [5] Ranzani OT, Prina E, Menéndez R, et al. New Sepsis Definition (Sepsis-3) and Community-acquired Pneumonia Mortality: A Validation and Clinical Decision-making Study [J]. Am J Respir Crit Care Med, 2017, 196(10): 1287-1297
- [6] Mangen MJ, Huijts SM, Bonten MJ, et al. The impact of community-acquired pneumonia on the health-related quality-of-life in elderly[J]. BMC Infect Dis, 2017, 17(1): 208-228
- [7] Viasus D, Núñez-Ramos JA, Vilorio SA, et al. Pharmacotherapy for community-acquired pneumonia in the elderly [J]. Expert Opin Pharmacother, 2017, 18(10): 957-964
- [8] Toledo D, Soldevila N, Torner N, et al. Factors associated with 30-day readmission after hospitalisation for community-acquired pneumonia in older patients: a cross-sectional study in seven Spanish regions[J]. Bmj Open, 2018, 8(3): e020243
- [9] Ceccato A, Cilloniz C, Ranzani OT, et al. Treatment with macrolides and glucocorticosteroids in severe community-acquired pneumonia: A post-hoc exploratory analysis of a randomized controlled trial [J]. Plos One, 2017, 12(6): e0178022
- [10] Cataudella E, Giraffa CM, Marca SD, et al. Neutrophil-To-Lymphocyte Ratio: An Emerging Marker Predicting Prognosis in Elderly Adults with Community-Acquired Pneumonia [J]. J Am Geriatr Soc, 2017, 65(8): 1796-1801
- [11] Takada T, Yamamoto Y, Terada K, et al. Diagnostic utility of appetite loss in addition to existing prediction models for community-acquired pneumonia in the elderly: a prospective diagnostic study in acute care hospitals in Japan[J]. Bmj Open, 2017, 7(11): e019155
- [12] Domínguez À, Soldevila N, Toledo D, et al. Effectiveness of 23-valent pneumococcal polysaccharide vaccination in preventing community-acquired pneumonia hospitalization and severe outcomes in the elderly in Spain[J]. Plos One, 2017, 12(2): e0171943
- [13] Min JC, Song JY, Ji YN, et al. Disease burden of hospitalized community-acquired pneumonia in South Korea: Analysis based on age and underlying medical conditions[J]. Medicine, 2017, 96(44): e8429
- [14] Saukkoriipi A, Palmu AA, Pascal T, et al. LytaQuantitative PCR on Sputum and Nasopharyngeal Swab Samples for Detection of Pneumococcal Pneumonia among the Elderly [J]. J Clin Microbiol, 2018, 56(1): e01231-e01217

- [15] Jwa H, Beom JW, Lee JH. Predictive Factors of Methicillin-Resistant *Staphylococcus aureus* Infection in Elderly Patients with Community-Onset Pneumonia[J]. *Tuberc Respir Dis*, 2017, 80(2): 201-209
- [16] Jokinen J, Snellman M, Palmu AA, et al. Testing Pneumonia Vaccines in the Elderly: Determining a Case Definition for Pneumococcal Pneumonia in the Absence of a Gold Standard [J]. *Am J Epidemiol*, 2018, 187(6): 1295-1302
- [17] Han X, Zhou F, Li H, et al. Effects of age, comorbidity and adherence to current antimicrobial guidelines on mortality in hospitalized elderly patients with community-acquired pneumonia [J]. *BMC Infect Dis*, 2018, 18(1): 192-214
- [18] Hariri G, Tankovic J, Boëlle PY, et al. Are third-generation cephalosporins unavoidable for empirical therapy of community-acquired pneumonia in adult patients who require ICU admission? A retrospective study[J]. *Ann Intensive Care*, 2017, 7(1): 35-50
- [19] Amalakuhan B, Echevarria KL, Restrepo MI. Managing community acquired pneumonia in the elderly - the next generation of pharmacotherapy on the horizon [J]. *Expert Opin Pharmacother*, 2017, 18(11): 1039-1048
- [20] Wongchew RM, García-León ML, Noyola DE, et al. Respiratory viruses detected in Mexican children younger than 5 years old with community acquired pneumonia. A national multicenter study[J]. *Int J Infect Dis*, 2017, 62(C): 32-38
- [21] Andersen SB, Baunbæk EG, Jensen AV, et al. Failure of CRP decline within three days of hospitalization is associated with poor prognosis of Community-acquired Pneumonia [J]. *Infect Dis*, 2017, 49(4): 251-260
- [22] Minnaard MC, Groot JAHD, Hopstaken RM, et al. The added value of C-reactive protein measurement in diagnosing pneumonia in primary care: a meta-analysis of individual patient data [J]. *CMAJ*, 2017, 189(2): E56-E63
- [23] Alcoba G, Keitel K, Maspoli V, et al. A three-step diagnosis of pediatric pneumonia at the emergency department using clinical predictors, C-reactive protein, and pneumococcal PCR [J]. *Eur J Pediatr*, 2017, 176(6): 815-824
- [24] Rodriguez GD, Rodriguez GD. A Novel Antimicrobial Stewardship Program-Guided Procalcitonin Initiative for Emergency Department Diagnosis of Bacterial Pneumonia in New York City[J]. *Open Forum Infect Dis*, 2017, 4(Suppl 1): S27-S28
- [25] Miroliaee AE, Salamzadeh J, Shokouhi S, et al. Effect of Vitamin D Supplementation on Procalcitonin as Prognostic Biomarker in Patients with Ventilator Associated Pneumonia Complicated with Vitamin D Deficiency[J]. *Iran J Pharm Res*, 2017, 16(3): 1254-1263
- [26] Masiá M, Padilla S, Ortiz d l TV, et al. Procalcitonin for selecting the antibiotic regimen in outpatients with low-risk community-acquired pneumonia using a rapid point-of-care testing: A single-arm clinical trial[J]. *Plos One*, 2017, 12(4): e0175634
- [27] Faluppecurariu OG, Diezdomingo J, Esposito S, et al. Clinical and laboratory features of children with community-acquired pneumonia are associated with distinct radiographic presentations [J]. *Eur J Pediatr*, 2018, 177(7): 1-10
- [28] Petersen PT, Egelund GB, Jensen AV, et al. Associations between biomarkers at discharge and co-morbidities and risk of readmission after community-acquired pneumonia: a retrospective cohort study[J]. *Eur J Clin Microbiol Infect Dis*, 2018, 1(3): 1103-1111
- [29] Pantzaris ND, Platanaki C, Pierrako C, et al. Neutrophil-to-lymphocyte Ratio Relation to Sepsis Severity Scores and Inflammatory Biomarkers in Patients with Community-acquired Pneumonia: A Case Series[J]. *J Transl Int Med*, 2018, 6(1): 43-46
- [30] Cannady SB, Hatten KM, Bur AM, et al. Use of free tissue transfer in head and neck cancer surgery and risk of overall and serious complication (s): An American College of Surgeons-National Surgical Quality Improvement Project analysis of free tissue transfer to the head and neck[J]. *Head Neck*, 2017, 39(4): 702-707
- [31] Shakeri F, Soukhtanloo M, Boskabady M H. The effect of hydro-ethanolic extract of *Curcuma longa* rhizome and curcumin on total and differential WBC and serum oxidant, antioxidant biomarkers in rat model of asthma [J]. *Iranian Journal of Basic Medical Sciences*, 2017, 20(2): e155
- [32] Benej M, Capov I, Skrickova J, et al. Association of the postoperative white blood cells (WBC) count in peripheral blood after radical surgical treatment of left upper lobe non-small cell lung cancer (NSCLC) with overall survival - single center results [J]. *Bratislavske Lekarske Listy*, 2017, 118(5): e299
- [33] Lu Y C, Wang C P, Yu T H, et al. Shift work is associated with metabolic syndrome in male steel workers-the role of resistin and WBC count-related metabolic derangements [J]. *Diabetology & Metabolic Syndrome*, 2017, 9(1): e83