

doi: 10.13241/j.cnki.pmb.2020.15.012

低频电刺激结合康复训练对脊髓损伤神经源性膀胱患者排尿症状、膀胱功能及生活质量的影响*

杨悦¹ 张治强² 杨颖¹ 蒋红英¹ 叶静¹ 杜春萍^{1Δ}

(1 四川大学华西医院康复医学中心 / 康复医学四川省重点实验室 四川 成都 610041;

2 成都高新海尔森医院康复科 四川 成都 610000)

摘要 目的:探讨低频电刺激结合康复训练对脊髓损伤(SCI)诱发神经源性膀胱(NB)患者排尿症状、膀胱功能及生活质量的影响。**方法:**选取2017年3月~2019年12月期间我院收治的SCI诱发NB患者97例,根据随机数字表法分为对照组(n=48)和研究组(n=49),对照组患者予以康复训练,研究组在对照组基础上联合低频电刺激,比较两组患者排尿症状、膀胱功能、生活质量及并发症发生情况。**结果:**两组治疗2周后、治疗4周后膀胱内压力、24 h单次排尿量、排尿次数、膀胱容量均较治疗前增加,且研究组高于对照组($P<0.05$);残余尿量均较治疗前下降,且研究组低于对照组($P<0.05$)。两组治疗2周后、治疗4周后膀胱功能积分均较治疗前下降,且研究组低于对照组($P<0.05$)。两组治疗4周后情感职能、躯体疼痛、精神健康、活力、生理职能、生理机能、健康状况、社会功能维度评分均升高,且研究组高于对照组($P<0.05$)。研究组并发症总发生率低于对照组($P<0.05$)。**结论:**低频电刺激结合康复训练治疗SCI诱发NB患者,可有效改善患者排尿症状,提高其膀胱功能及生活质量,同时还可减少并发症发生率,具有一定的临床应用价值。

关键词:低频电刺激;康复训练;脊髓损伤;神经源性膀胱;排尿症状

中图分类号:R651.2;R694.6;Q64 **文献标识码:**A **文章编号:**1673-6273(2020)15-2863-05

Effects of Low Frequency Electrical Stimulation Combined with Rehabilitation Training on Micturition Symptoms, Bladder Function and Quality of Life of Neurogenic Bladder Patients with Spinal Cord Injury*

YANG Yue¹, ZHANG Zhi-qiang², YANG Ying¹, JIANG Hong-ying¹, YE Jing¹, DU Chun-ping^{1Δ}

(1 Rehabilitation Medical Center/Sichuan Key Laboratory of Rehabilitation Medicine,

West China Hospital of Sichuan University, Chengdu, Sichuan, 610041, China;

2 Department of Rehabilitation, Chengdu High Tech Haiersan Hospital, Chengdu, Sichuan, 610000, China)

ABSTRACT Objective: To investigate the effect of low frequency electrical stimulation combined with rehabilitation training on micturition symptoms, bladder function and quality of life of neurogenic bladder (NB) patients induced by spinal cord injury (SCI). **Methods:** 97 patients with NB induced by SCI in our hospital from March 2017 to December 2019 were selected, patients were divided into control group (n=48) and study group (n=49) according to the method of random number table. The patients in the control group were given rehabilitation training. The study group was combined with low frequency electrical stimulation on the basis of the control group. The urination symptoms, bladder function, quality of life and complications of the two groups were compared. **Results:** 2 weeks and 4 weeks after treatment, the number of Intravesical pressure, 24 h single micturition volume, times of micturition, bladder volume in the two groups increased compared with before treatment, and study group was higher than control group ($P<0.05$). The residual urine volume was lower than that before treatment, study group was lower than control group ($P<0.05$). 2 weeks and 4 weeks after treatment, the bladder function scores of the two groups were lower than before treatment, study group was lower than control group ($P<0.05$). The scores of emotional function, physical pain, mental health, vitality, physiological function, physiological function, health status and social function in the two groups increased, study group were higher than control group ($P<0.05$). The total incidence of complications in the study group was lower than that in the control group ($P<0.05$). **Conclusion:** Low frequency electrical stimulation combined with rehabilitation training can effectively improve the symptoms of micturition, improve the bladder function and quality of life of patients with NB induced by SCI, at the same time, and it can reduce the incidence of complications, and have a high clinical application value.

Key words: Low frequency electrical stimulation; Rehabilitation training; Spinal cord injury; Neurogenic bladder; Micturition symptoms

Chinese Library Classification(CLC): R651.2; R694.6; Q64 **Document code:** A

Article ID: 1673-6273(2020)15-2863-05

* 基金项目:国家自然科学基金项目(81572231)

作者简介:杨悦(1990-),女,硕士研究生,研究方向:康复医学,E-mail: yy5201533@163.com

Δ 通讯作者:杜春萍(1971-),女,本科,教授,研究方向:康复医学,E-mail: ducp118@163.com

(收稿日期:2020-03-04 接受日期:2020-03-27)

前言

神经源性膀胱(NB)是指控制排尿功能的中枢神经系统或周围神经受到损害,进而引起膀胱尿道功能障碍的一种疾病^[1]。脊髓损伤(SCI)是引起NB的主要原因之一,当机体神经环路损伤时,易引发NB,产生尿潴留及尿不畅等症状,若未能及时予以治疗,可诱发泌尿系统并发症,如肾衰竭、上尿路损害等,严重者甚至导致患者死亡^[2,3]。现临床有关SCI诱发NB的治疗方案尚未完全统一,主要集中于药物、外科手术及膀胱功能康复训练等,其中药物治疗存在不良反应大的不足,外科手术则会对机体造成创伤,因此临床中多以膀胱功能康复训练为主,但仍疗效一般,无法达到预期的治疗效果^[4-6]。低频电刺激是将超低脉宽的低频电流输入人体靶向治疗点,继而促进神经系统功能重建以改善患者症状的一种治疗方式^[7-9]。本研究通过对我院收治的部分SCI诱发NB患者给予低频电刺激结合康复训练治疗,取得了较好的疗效,现整理报道如下。

1 资料与方法

1.1 一般资料

纳入标准:(1)经临床表现、影像学检查确诊为SCI;(2)伴有NB症状;(3)患者及其家属知情本研究且签署了同意书;(4)患者意识清晰,生命体征平稳。排除标准:(1)既往行膀胱造漏术手术损伤者;(2)意识不清或有精神疾病无法配合治疗者;(3)因前列腺增生或其他疾病导致排尿障碍;(4)心脑血管等重要脏器严重疾病者。选取2017年3月~2019年12月期间我院收治的SCI诱发NB患者97例,根据随机数字表法分为对照组(n=48)和研究组(n=49),其中研究组男27例,女22例,年龄25~50岁,平均(43.19±4.25)岁;损伤节段包括颈椎20例,胸椎15例,腰椎14例;SCI病程3~10月,平均(6.57±0.96)月。对照组男28例,女20例,年龄27~53岁,平均(42.91±4.67)岁;损伤节段包括颈椎19例,胸椎16例,腰椎13例;SCI病程2~9月,平均(6.29±0.87)月。两组患者一般资料对比未见统计学差异($P>0.05$),组间具有可比价值。本次研究已通过我院伦理学委员会批准进行。

1.2 方法

两组均接受膀胱康复训练治疗,具体如下:(1)规律的饮水计划:要求患者晨起至20:00每隔2小时饮水200~250 mL,期间若进食流质、粥、汤、果汁等,应减去相应分量的水分。平均每天总入液量控制为1500-2000 mL左右。(2)清洁间歇导尿,为患者建立膀胱充盈排泄规律。(3)盆底肌功能训练。指导患者在立、卧或站位下进行尿道和肛门主动收缩训练,收缩与放松持续时间分别5~10s,20组/次,每日5次。(4)诱导排尿训练:a.利用条件反射诱导排尿。可以离床患者,协助患者坐于坐厕,让患者听流水声。需要卧床患者,放置便器,用温热毛巾外敷膀胱区域或者用温水冲洗会阴部,边冲边轻轻按摩患者膀胱膨隆区域。b.开塞露塞肛诱导排尿。采用开塞露塞肛,促使逼尿肌收缩,内括约肌松弛从而达到排尿。(5)肛门括约肌训练:指导患者主动进行肛门牵拉的训练,医务人员使用中指或食指两个指节,戴上指套润滑后,分别朝3点、6点、9点、12点四个方向进行牵拉,促使盆底肌、尿道括约肌痉挛缓解,每次牵拉收缩持续

时间控制在10s左右,训练10次为1组,每日5组。上述膀胱康复训练持续治疗4周。研究组在膀胱康复训练的基础上联合低频电刺激治疗,具体如下:选择深圳市力合医疗器械有限公司生产的LGT-1000B低频电子脉冲膀胱治疗仪,其中膀胱近顶部侧壁贴两个电极片,耻骨、肚脐连线膀胱顶下缘及骶尾关节处贴两个电极片,接通电源。电刺激期间,设置为患者能耐受的最大电流强度,以适量肌肉收缩为宜。1次/d,4次/周,持续治疗4周。

1.3 观察指标

(1)记录两组患者治疗前、治疗2周后、治疗4周后的排尿症状指标,包括:膀胱内压力、日排尿次数、24h单次排尿量、残余尿量、膀胱容量。其中残余尿量及膀胱容量均通过膀胱容量测定仪测量;膀胱内压力采用尿流动力学检查评定。24h单次排尿量通过收集两组患者24h尿量,取平均值测定。(2)记录两组患者治疗前、治疗2周后、治疗4周后的膀胱功能积分。其中0分:通过反射可自行排尿,排尿及终止排尿受意识控制,残余尿量<50 mL;1分:通过反射刺激能自行排尿,排尿及终止排尿缓慢但受意识控制,残余尿量为50~150 mL;2分:经反射刺激能排尿,排尿及终止排尿不完全受意识控制,残余尿量150~250 mL;3分:不能通过反射刺激自行排尿,排尿及终止排尿不受意识控制,膀胱容量低于400 mL^[10]。(3)采用生活质量调查简表(SF-36)^[11]评价患者治疗前、治疗4周后的生活质量,其中SF-36包括情感职能、躯体疼痛、精神健康、活力、生理职能、生理机能、健康状况、社会功能这8个维度,每个维度均为100分,分数越高,生活质量越好。(4)记录两组患者治疗4周后的肾功能损害发生率、尿路感染发生率及泌尿系统造影异常发生率。

1.4 统计学方法

数据分析采用SPSS24.0软件进行,计量资料以($\bar{x} \pm s$)表示,实施t检验,计数资料以[n(%)]表示,实施 χ^2 检验,检验标准设置为 $\alpha=0.05$ 。

2 结果

2.1 两组排尿症状比较

两组治疗前日排尿次数、残余尿量、24h单次排尿量、膀胱内压力、膀胱容量比较差异均无统计学意义($P>0.05$);两组治疗2周后、治疗4周后日膀胱内压力、24h单次排尿量、排尿次数、膀胱容量均较治疗前增加,且研究组高于对照组($P<0.05$);残余尿量均较治疗前降低,且研究组低于对照组($P<0.05$);详见表1。

2.2 膀胱功能比较

两组治疗前膀胱功能积分比较无差异($P>0.05$);两组治疗2周后、治疗4周后膀胱功能积分均较治疗前下降,且研究组低于对照组($P<0.05$);详见表2。

2.3 两组生活质量比较

两组治疗前情感职能、躯体疼痛、精神健康、活力、生理职能、生理机能、健康状况、社会功能维度评分比较差异无统计学意义($P>0.05$);两组治疗4周后情感职能、躯体疼痛、精神健康、活力、生理职能、生理机能、健康状况、社会功能维度评分均升高,且研究组高于对照组($P<0.05$);详见表3。

表 1 两组排尿症状比较($\bar{x} \pm s$)Table 1 Comparison of micturition symptoms between the two groups($\bar{x} \pm s$)

Groups	Time	Number of daily micturition(n/d)	Residual urine volume(mL)	24 hours of single micturition(ml)	Bladder pressure (mmH ₂ O)	Bladder volume (mL)
Control group (n=48)	Before treatment	2.26±0.31	291.74±16.82	46.12±7.61	12.92±2.98	329.15±16.12
	2 weeks after treatment	3.45±0.52 ^a	262.76±17.73 ^a	73.44±7.66 ^a	16.06±2.48 ^a	368.25±17.48 ^a
	4 weeks after treatment	4.96±0.34 ^{ab}	237.39±20.27 ^{ab}	91.76±10.23 ^{ab}	20.18±2.31 ^{ab}	394.97±20.64 ^{ab}
Study group(n=49)	Before treatment	2.21±0.29	289.85±18.33	45.79±6.29	12.62±2.27	328.04±18.36
	2 weeks after treatment	4.87±0.38 ^{ac}	234.38±17.06 ^{ac}	99.74±7.82 ^{ac}	22.81±2.76 ^{ac}	402.67±17.73 ^{ac}
	4 weeks after treatment	6.79±0.43 ^{abc}	202.76±16.72 ^{abc}	134.68±12.79 ^{abc}	26.45±2.52 ^{abc}	454.73±16.64 ^{abc}

表 2 两组膀胱功能比较($\bar{x} \pm s$, 分)Table 2 Comparison of bladder function between the two groups($\bar{x} \pm s$, scores)

Groups	Before treatment	2 weeks after treatment	4 weeks after treatment
Control group(n=48)	2.97±0.26	2.35±0.21 ^a	1.78±0.19 ^{ab}
Study group(n=49)	2.92±0.21	1.86±0.24 ^a	1.23±0.14 ^{ab}
t	1.043	10.693	16.254
P	0.300	0.000	0.000

Note: compared with before treatment, ^a $P<0.05$; compared with 2 weeks after treatment, ^b $P<0.05$.

表 3 两组生活质量比较($\bar{x} \pm s$, 分)Table 3 Comparison of quality of life between the two groups($\bar{x} \pm s$, scores)

Groups	Time	Emotional function	Physical pain	Mental health	Vitality	Physiological function	Physiological function	Health status	Social function
Control group(n=48)	Before treatment	46.89±6.30	53.27±7.24	51.27±7.24	55.37±6.83	51.27±8.26	55.73±6.46	54.58±7.32	50.28±6.86
	4 weeks after treatment	71.09±7.27 ^a	73.19±7.75 ^a	74.13±6.26 ^a	72.57±7.32 ^a	74.31±7.16 ^a	72.87±7.23 ^a	76.05±6.25 ^a	72.47±8.21 ^a
Study group (n=49)	Before treatment	46.26±7.41	53.66±7.34	50.93±8.68	54.95±7.92	50.96±7.73	55.44±7.66	55.06±7.48	50.75±7.48
	4 weeks after treatment	85.14±7.13 ^{ab}	83.18±7.73 ^{ab}	84.92±7.62 ^{ab}	88.28±7.34 ^{ab}	86.54±7.25 ^{ab}	85.14±7.35 ^{ab}	86.95±7.48 ^{ab}	88.37±8.61 ^{ab}

Note: compared with before treatment, ^a $P<0.05$; compared with control group, ^b $P<0.05$.

2.4 并发症发生情况比较

研究组并发症总发生率低于对照组($P<0.05$);详见表 4。

3 讨论

近年来,随着各种医疗条件的改善和医疗技术的进步,临床中针对 SCI 患者的治疗效果已经有了显著提升,但绝大多数 SCI 患者仍无法彻底治愈,常存在感觉、运动及自主神经等功能障碍,SCI 造成的影响可伴随终身,严重降低其生活质量^[12-14]。

表 4 两组并发症发生情况比较例(%)

Table 4 Comparison of complications between the two groups n(%)

Groups	Urinary tract infection	Renal function damage	Abnormal urography	Total incidence rate
Control group(n=48)	4(8.33)	6(12.50)	5(10.42)	15(31.25)
Study group(n=49)	2(4.08)	2(4.08)	1(2.04)	5(10.20)
χ^2			5.562	
P			0.010	

NB 作为 SCI 的主要并发症之一, 会使患者因膀胱功能障碍而表现出异常的排尿症状, 其发生原因可能在于尿潴留造成膀胱充盈膨胀, 造成黏膜充血、水肿, 加上膀胱内尿液滞留时间过长, 导致细菌大量滋生、繁殖, 增加尿路感染发生风险, 进而诱发 NB^[15-17]。此外, SCI 诱发 NB 后还可引起败血症、肾功能不全等, 加重病情, 甚至导致患者死亡。国内外的研究资料均表明^[18,19], SCI 诱发 NB 患者的膀胱功能及症状变化均是影响其治疗效果和生存质量的主要因素。针对 SCI 诱发 NB 患者的主要治疗目标为改善临床症状、保护肾功能、减少上尿路的损害、改善膀胱功能, 并通过减少并发症来提高其生活质量^[20]。既往常规的治疗方法是膀胱功能训练结合间歇导尿, 但是这种方法必须将导尿管长期留置于膀胱内, 易增加并发症的发生率, 最终致使膀胱的容量减小, 疗效不甚理想^[21]。而低频电刺激治疗对感觉、运动神经都有强的刺激作用, 近年来已逐渐应用于 SCI 诱发 NB 的辅助治疗中^[22]。

本研究结果显示, 低频电刺激结合康复训练治疗 SCI 诱发 NB 患者, 可有效改善患者排尿症状及膀胱功能。其作用机制可能如下: 低频电刺激可解除脊髓排尿中枢的抑制, 诱导排尿反射, 使逼尿肌持续收缩, 尿道括约肌扩张, 进而促进尿液排出, 降低膀胱内压力, 减少膀胱内残余尿量^[23,24]。此外, 低频电刺激可兴奋膀胱神经, 进而有效改善膀胱局部的血液循环, 促进膀胱排尿功能恢复^[25]。当机体膀胱功能正常时, 其具备低压储存和意愿排泄这两方面的功能, 同时呈周期性变换。而低频电刺激具有选择性激活骶神经根中支配膀胱逼尿肌的直径较小副交感神经的作用, 骶神经根主要包含支配尿道括约肌的粗大神经纤维和支配膀胱逼尿肌的细小神经纤维以及下肢肌肉的神经纤维, 低频电刺激时可先激活粗大神经纤维, 其次再激活细胞的神经纤维, 故而在治疗时, 尿道括约肌总是先于膀胱逼尿肌收缩, 导致前者压力高于后者, 膀胱无法排除尿液。而当电刺激结束后, 尿道压力立即下降, 而膀胱压力可缓慢下降, 在这段时间内排尿可刺激膀胱功能恢复, 从而改善排尿症状^[26-28]。此外, 两组患者生活质量均有所改善, 且低频电刺激结合康复训练治疗者的改善效果更佳。推测其原因为低频电刺激具有安全、有效、无创、廉价等优势, 且治疗期间设定多种安全装置: 过流保护、短路保护、过压保护、输出强度自动锁定装置, 可更为精确的控制对患者的电刺激, 促进其功能恢复。治疗过后患者症状得到显著改善, 有效促进膀胱反射性机制的建立, 使疾病对患者日常生活及功能的影响随着病情的缓解而减轻, 同时还能树立起患者战胜疾病的信心^[29]。另外, 低频电刺激结合康复训练治疗可有效减少 SCI 诱发 NB 患者的并发症, 这主要是因为低频电刺激结合康复训练治疗可有效缓解患者的各种症状, 减少机体组织受到的压迫且可改善新陈代谢、减轻受损组织水肿等, 最终推进受损部位循环修复, 并进一步减少并发症发生风险^[30]。

综上所述, 低频电刺激结合康复训练治疗 SCI 诱发 NB 患者, 可有效改善患者排尿症状, 提高其膀胱功能及生活质量, 同时还可减少并发症发生率, 具有一定的临床应用价值。

参考文献(References)

- [1] Erdogru T, Onur R. Laparoscopic selective pudendal neuromodulation for neurogenic bladder: a review of literature [J]. Cent European J Urol, 2020, 73(1): 39-41
- [2] Doherty SP, Vanhoostenberghe A, Duffell LD, et al. Ambulatory urodynamic monitoring assessment of dorsal genital nerve stimulation for suppression of involuntary detrusor contractions following spinal cord injury: a pilot study[J]. Spinal Cord Ser Cases, 2020, 6(1): 30
- [3] Tien NLB, Hoa ND, Thanh VV, et al. utologous Transplantation of Adipose-Derived Stem Cells to Treat Acute Spinal Cord Injury: Evaluation of Clinical Signs, Mental Signs, and Quality of Life[J]. Open Access Maced J Med Sci, 2019, 7(24): 4399-4405
- [4] Welk B, Lenherr S, Elliott S, et al. The creation and validation of a short form of the Neurogenic Bladder Symptom Score [J]. NeurourolUrodyn, 2020, 39(4): 1162-1169
- [5] Tate DG, Wheeler T, Lane GI, et al. Recommendations for evaluation of neurogenic bladder and bowel dysfunction after spinal cord injury and/or disease[J]. J Spinal Cord Med, 2020, 43(2): 141-164
- [6] Gater DR. Neurogenic bowel and bladder evaluation strategies in spinal cord injury: New directions[J]. J Spinal Cord Med, 2020, 43(2): 139-140
- [7] Cakan C, Obermayer K. Biophysically grounded mean-field models of neural populations under electrical stimulation [J]. PLoSComputBiol, 2020, 16(4): e1007822
- [8] Helbing C, Angenstein F. Frequency-dependent electrical stimulation of fimbria-fornix preferentially affects the mesolimbic dopamine system or prefrontal cortex[J]. Brain Stimul, 2020, 13(3): 753-764
- [9] Balderston NL, Beydler EM, Goodwin M, et al. Low-frequency parietal repetitive transcranial magnetic stimulation reduces fear and anxiety [J]. Transl Psychiatry, 2020, 10(1): 68
- [10] 姜天伟, 田梦, 高振梅, 等. 温针结合膀胱功能训练治疗脊髓损伤后神经源性膀胱的临床研究 [J]. 浙江中医药大学学报, 2017, 41(10): 840-844
- [11] 彭颖, 陈露, 卓琳, 等. SF-36 和 EQ-5D 量表在慢性肾脏病患者生命质量评估中的对比研究[J]. 东南大学学报(医学版), 2019, 38(6): 1024-1028
- [12] Abolhasanpour N, Hajebrahimi S, Ebrahimi-Kalan A, et al. Urodynamic Parameters in Spinal Cord Injury-Induced Neurogenic Bladder Rats after Stem Cell Transplantation: A Narrative Review[J]. Iran J Med Sci, 2020, 45(1): 2-15
- [13] Zhang Y, Li L, Mu J, et al. Implantation of a functional TEMPO-hydrogel induces recovery from rat spinal cord transection through promoting nerve regeneration and protecting bladder tissue [J]. BiomaterSci, 2020, 8(6): 1695-1701
- [14] Schaeffer EM. Re: Bladder Emptying Method is the Primary Determinant of Urinary Tract Infections in Patients with Spinal Cord Injury: Results from a Prospective Rehabilitation Cohort Study [J]. J Urol, 2020, 203(4): 644
- [15] Lei H, Fu Y, Xu G, et al. Different types of acupuncture and moxibustion therapy for neurogenic bladder after spinal cord injury: A systematic review and network meta-analysis study protocol [J]. Medicine (Baltimore), 2020, 99(1): e18558
- [16] 王小雪, 张健, 丛双, 等. 针刺联合生物反馈治疗脊髓损伤后神经源性膀胱的疗效观察[J]. 现代生物医学进展, 2018, 18(22): 4303-4307
- [17] Kinnear N, Barnett D, O'Callaghan M, et al. The impact of catheter-based bladder drainage method on urinary tract infection risk in spinal

- cord injury and neurogenic bladder: A systematic review[J]. *Neurourol Urodyn*, 2020, 39(2): 854-862
- [18] 王能, 冯琦钒, 林晓伟, 等. 针灸联合金匱肾气丸对脊髓损伤后神经源性膀胱尿潴留患者影响的临床研究[J]. *世界中西医结合杂志*, 2020, 15(3): 524-527, 531
- [19] Daia C, Bumbea AM, Badiu CD, et al. Interferential electrical stimulation for improved bladder management following spinal cord injury[J]. *Biomed Rep*, 2019, 11(3): 115-122
- [20] Elliott CS, Stoffel JT, Myers JB, et al. Validation of Upper Extremity Motor Function as a Key Predictor of Bladder Management After Spinal Cord Injury[J]. *Arch Phys Med Rehabil*, 2019, 100(10): 1939-1944
- [21] Fang H, Lin J, Liang L, et al. A nonsurgical and nonpharmacological care bundle for preventing upper urinary tract damage in patients with spinal cord injury and neurogenic bladder[J]. *Int J Nurs Pract*, 2020, 26(2): e12761
- [22] Irahara T, Sato N, Otake K, et al. Neuromuscular Electrical Stimulation Improves Energy Substrate Metabolism and Survival in Mice With Acute Endotoxic Shock[J]. *Shock*, 2020, 53(2): 236-241
- [23] Setya A, Nair P, Cheng SX. Gastric electrical stimulation: An emerging therapy for children with intractable gastroparesis[J]. *World J Gastroenterol*, 2019, 25(48): 6880-6889
- [24] Couturier NH, Durand DM. Comparison of fiber tract low frequency stimulation to focal and ANT stimulation in an acute rat model of focal cortical seizures[J]. *Brain Stimul*, 2020, 13(2): 499-506
- [25] Dillon MT, Buss E, O'Connell BP, et al. Low-Frequency Hearing Preservation With Long Electrode Arrays: Inclusion of Unaided Hearing Threshold Assessment in the Postoperative Test Battery[J]. *Am J Audiol*, 2020, 29(1): 1-5
- [26] Su S, Chai G, Shu X, et al. Electrical stimulation-induced SSSEP as an objective index to evaluate the difference of tactile acuity between the left and right hand[J]. *J Neural Eng*, 2020, 17(1): 016053
- [27] Koshy SM, Wiesman AI, Spooner RK, et al. Multielectrode Transcranial Electrical Stimulation of the Left and Right Prefrontal Cortices Differentially Impacts Verbal Working Memory Neural Circuitry[J]. *Cereb Cortex*, 2020, 30(4): 2389-2400
- [28] Wakahara T, Shiraogawa A. Effects of neuromuscular electrical stimulation training on muscle size in collegiate track and field athletes[J]. *PLoS One*, 2019, 14(11): e0224881
- [29] Zeng FG, Tran P, Richardson M, et al. Human Sensation of Transcranial Electric Stimulation[J]. *Sci Rep*, 2019, 9(1): 15247
- [30] 郝丽霞, 张琰, 刘帅, 等. 低频电刺激结合康复训练治疗脊髓损伤神经源性膀胱患者疗效观察[J]. *中国临床研究*, 2017, 30(1): 116-119

(上接第 2837 页)

- [20] Adam Bryant, Catalina A Palma, Vivek Jayaswal, et al. miR-10a is aberrantly overexpressed in Nucleophosmin1 mutated acute myeloid leukaemia and its suppression induces cell death[J]. *Mol Cncr*, 2012, 11: 8
- [21] Dong J, Zhang Z, Huang H, et al. miR-10a rejuvenates aged human mesenchymal stem cells and improves heart function after myocardial infarction through KLF4 [J]. *Stem Cell Research & Therapy*, 2018, 9(1): 151
- [22] Xue H, Tian G Y. MiR-429 regulates the metastasis and EMT of HCC cells through targeting RAB23 [J]. *Archives of Biochemistry & Biophysics*, 2017, 637: 48
- [23] Xu Z, Gakhar L, Bain F E, et al. The Tiam1 guanine nucleotide exchange factor is auto-inhibited by its pleckstrin homology coiled-coil extension domain[J]. *Journal of Biological Chemistry*, 2017, 292
- [24] VAV2, a guanine nucleotide exchange factor for Rac1, regulates glucose-stimulated insulin secretion in pancreatic beta cells [J]. *Diabetologia*, 2015, 58(11): 2573-2581
- [25] Meghan E. Minard, Lee-Su Kim, Janet E. Price, et al. The Role of the Guanine Nucleotide Exchange Factor Tiam1 in Cellular Migration, Invasion, Adhesion and Tumor Progression [J]. *Breast Cancer Research & Treatment*, 2004, 84(1): 21-32
- [26] Chen G, Lu L, Liu C, et al. MicroRNA-377 Suppresses Cell Proliferation and Invasion by Inhibiting TIAM1 Expression in Hepatocellular Carcinoma[J]. *PLOS ONE*, 2015, 10(3): e0117714
- [27] Izumi D, Toden S, Ureta E, et al. TIAM1 promotes chemoresistance and tumor invasiveness in colorectal cancer [J]. *Cell Death and Disease*, 2019, 10(4): 267
- [28] Jia L, Li Z, Liu Q, Piao J, et al. Clinicopathological implications of Tiam1 overexpression in invasive ductal carcinoma of the breast[J]. *BMC Cancer*, 2016, 16(1): 681
- [29] Han W, Wang L, Zhang L, et al. Circular RNA circ-RAD23B promotes cell growth and invasion by miR-593-3p/CCND2 and miR-653-5p/TIAM1 pathways in non-small cell lung cancer [J]. *Biochemical and Biophysical Research Communications*, 2019, 510(3): 462-466
- [30] Wang J X, Zhou Y N, Zou S J, et al. Expression of Tiam1 in gastric cancer tissue and its relationship with clinicopathologic characteristics and survival time[J]. *China Practical Medicine*, 2015, 10(32)