

doi: 10.13241/j.cnki.pmb.2021.14.042

· 专论与综述 ·

外泌体在淋巴瘤中的研究进展*

陈艳¹ 梁奇² 杨志武² 刘珊² 陈文^{3Δ}

(益阳市中心医院 1 中西医结合科; 2 血液科 湖南 益阳 413000; 3 解放军总医院第八医学中心 北京 100091)

摘要: 病毒感染和环境污染等使得淋巴瘤的发病率逐年升高, 早期诊断和精准治疗具有十分重要的临床意义。外泌体是一种脂质双层膜结构的微小囊泡, 介导了细胞间交流和信息交换。近几年许多研究证实外泌体是淋巴瘤的发生、进展和耐药的重要机制。外泌体内核酸和小分子可用于淋巴瘤的早期诊断和预测患者预后。材料学修饰可显著增强外泌体治疗的靶向性和治疗效能。本文总结了外泌体生物学特性、分离和鉴定方法、与肿瘤相关性、及其在淋巴瘤中的研究进展, 为淋巴瘤的预警和治疗提供参考。

关键词: 淋巴瘤; 外泌体; 早期诊断; 精准治疗

中图分类号: R733.4 **文献标识码:** A **文章编号:** 1673-6273(2021)14-2797-04

Research Progress of Exosomes in Lymphoma*

CHEN Yan¹, LIANG Qi², YANG Zhi-wu², LIU Shan², CHEN Wen^{3Δ}

(1 Department of Integrated TCM & Western Medicine; 2 Department of Hematology, Yiyang Central Hospital, Yiyang, Hunan, 413200, China; 3 The 8th medical center of PLA General Hospital, Beijing, 100091, China)

ABSTRACT: Viral infection and environmental pollution make the incidence of lymphoma increase year by year. Early diagnosis and precise treatment are of great clinical significance. Exosomes are tiny vesicles of lipid bilayer membrane structure, which mediate cell to cell communication and information exchange. In recent years, many studies have proved that exosomes are the important mechanism of the occurrence, development and drug resistance of lymphoma. Nucleic acids and small molecules in exosomes can be used for early diagnosis and prognosis of lymphoma. Material modification can significantly enhance the targeting and therapeutic efficacy of exosomes. This paper summarized the biological characteristics of exosomes, the methods of isolation and identification, the relationship between exosomes and tumor, and the research progress of exosomes in lymphoma, to provide references for the early warning and treatment of lymphoma.

Key words: Lymphoma; Exosomes; Early diagnosis; Precise treatment

Chinese Library Classification(CLC): R733.4 **Document code:** A

Article ID: 1673-6273(2021)14-2797-04

前言

近几年淋巴瘤的发病呈快速上升和年轻化趋势, 已经成为血液系统最常见的恶性肿瘤, 可分为非霍奇金淋巴瘤(non-Hodgkin's lymphoma, NHL) 和霍奇金淋巴瘤(Hodgkin's lymphoma, HL)两大类^[1]。淋巴瘤症状包括发热、无痛性淋巴结肿大和全身器官组织受累症状等, 大部分不具有临床特异性, 早期诊断十分困难, 很多患者发现时已经进入了晚期^[2]。化疗和放疗是淋巴瘤的主要治疗手段, 可以获得暂时较好的缓解, 但是淋巴瘤的复发率十分高, 统计数据显示约有 40% 的弥漫性大 B 细胞淋巴瘤在治疗后数月或数年间会复发, 且复发后往往是很难治愈的^[3]。因此寻找淋巴瘤诊断和治疗的新靶点具有十分重要临床意义。大量研究证实外泌体(exosome)是淋巴瘤发生

和进展的重要机制, 可用于淋巴瘤的早期诊断和靶向治疗^[4]。本文总结了外泌体生物学特征及其在淋巴瘤中的最新进展, 为其诊断和治疗提供理论参考。

1 外泌体的生物学特征

外泌体是一种细胞分泌具有脂质双层膜结构的微小囊泡, 直径约为 30-120 nm, 内含丰富的脂类、核酸和蛋白质等小分子, 是细胞间交流和信息交换的重要机制。外泌体表达生物膜特异性标志, 包括 CD63、CD9、CD81 和肿瘤敏感基因 101(tumor sensitive gene 101, TSG101)等^[5]。此外外泌体还保留着供体细胞特征, 高表达相应的表面标志, 因此通过检测这些蛋白的表达可以追踪外泌体的来源^[6]。

目前认为外泌体生物学功能的发挥主要通过三条途径:

* 基金项目: 湖南省科技创新计划项目(2018SK52204); 国家自然科学基金青年基金项目(31800814)

作者简介: 陈艳(1983-), 女, 医学硕士, 副主任医师, 主要研究方向: 肿瘤的中西医结合治疗

Δ 通讯作者: 陈文(1987-), 男, 医学博士, 副主任医师, 硕士研究生导师, 主要研究方向: 肿瘤的靶向治疗,

电话: 13752985404, E-mail: 1607283686@qq.com

(收稿日期: 2020-12-23 接受日期: 2021-01-18)

结合靶细胞细胞膜,融合成为新的细胞表面分子;结合靶细胞表面受体,激活细胞内特定的信号通路;与细胞融合或者被消化后,释放内含蛋白质或者核酸小分子,调控靶细胞功能^[7]。

2 外泌体的分离和鉴定

去除细胞碎片或者蛋白质影响,富集到合适的浓度是外泌体分离的关键。目前常用的外泌体分离方法包括差速超速离心法、聚乙二醇(polyethylene glycol,PEG)沉淀法、多孔结构诱捕法、免疫磁珠法和沉淀试剂盒法等。目前应用较多的为沉淀试剂盒法、免疫磁珠法和差速超速离心法^[8]。每种分离方法均有各自的优点和不足之处,研究者应该根据自己需求选择最合适的方法。

2006年,外泌体典型的形态结构图第一次发布,之后科学家逐渐制定了外泌体鉴定的标准和常用方法。透射电镜下呈茶托型或一侧凹陷的半球形的双层囊膜超微结构,直径30-120 nm^[9];粒径分析仪检测外泌体大小及粒径分布;Western blot 或者流式细胞术检测外泌体特异性表面标志,比如TSG101、CD9、CD81和CD63等。另外一些指标比如Alix、Syntenin和热休克蛋白70(heat shock protein 70,HSP70)也有一定的辅助鉴定作用^[10]。这些监测指标均存在不足之处,比如透射电镜只关注少量外泌体特征,很难检测整个样本的纯度;外泌体特异性表面标志检测缺乏合适的阳性或者阴性对照。

3 外泌体与恶性肿瘤

外泌体的生物学功能与供体细胞紧密相关,因此对肿瘤具有双重作用。T细胞、树突细胞、自然杀伤细胞(natural killer cell,NK细胞)和巨噬细胞源性外泌体内含IL-6和IL-1等免疫活性分子,在肿瘤内可激活免疫反应,抑制肿瘤生长和转移^[11-14]。免疫抑制细胞,比如间充质干细胞、调节性T细胞(regulatory T cell,Treg细胞)和M2型巨噬细胞源性外泌体则高表达细胞毒性T淋巴细胞相关抗原-4(cytotoxic T lymphocyte associated antigen-4,CTLA-4)、人白细胞抗原-G(human leucocyte antigen-G,HLA-G)和糖皮质激素诱导的肿瘤坏死因子受体(glucocorticoid-induced tumor necrosis factor receptor,GITR)等免疫抑制分子,能够诱导T细胞凋亡,抑制抗肿瘤免疫反应,促进肿瘤增殖、侵袭和转移^[15-17]。此外,外泌体还能通过其他方式发挥生物学功能,比如内皮祖细胞(endothelial progenitor cell,EPC)源性外泌体内含血管内皮生长因子(vascular endothelial growth factor,VEGF)、IL-10和基质金属蛋白酶9(matrix metalloproteinase 9,MMP9)等,可诱导肿瘤生长、血管生成、周围组织浸润和远处转移等^[18-21]。

恶性肿瘤内部缺血缺氧微环境使得肿瘤细胞能够分泌更多的外泌体。肿瘤细胞分泌的外泌体在一定程度上可激活免疫反应,发挥抗肿瘤功能。但是这种效果是十分局限和微弱的,总的来说,肿瘤细胞源性外泌体主要发挥免疫抑制和促肿瘤进展的作用^[7,22,23]。胃癌细胞源性外泌体内含大量转化生长因子- β (transforming growth factor- β ,TGF- β)和IL-10,可增加淋巴细胞分类中Treg细胞比例,诱导T细胞和NK细胞凋亡,抑制抗肿瘤免疫反应。此外,胃癌细胞分泌的外泌体还能激活丝裂原活化蛋白激酶(mitogen-activated protein kinase,MAPK)、细胞

外调节蛋白激酶(extracellular regulated protein kinase,ERK)等信号通路,直接诱导胃癌的生长、浸润和转移,CD97在这些过程中扮演着关键角色^[10,24,25]。外泌体与化疗、靶向治疗耐药也具有很大的相关性,许多研究显示耐药癌细胞分泌的外泌体可将一些microRNA(miR)、LncRNA和小分子靶向递送给敏感细胞,促进敏感癌细胞对曲妥珠单抗、替莫唑胺、伊马替尼和紫杉醇的耐药^[26-28]。

4 外泌体与淋巴瘤

外泌体在淋巴瘤进展、复发和耐药中均发挥着重要的作用。免疫细胞源性外泌体可进入淋巴瘤瘤体内,具有一定抗肿瘤功能。淋巴瘤源性外泌体可被树突细胞摄取,在一定程度上能激活免疫反应^[29,30]。但是对于淋巴瘤患者,这些作用往往是十分有限的。很多细胞分泌的外泌体发挥着促进淋巴瘤进展的作用。内皮细胞或者EPC源性外泌体可促进淋巴瘤血管生成;Treg在淋巴瘤患者外周血淋巴细胞分类中比例显著升高,分泌的外泌体可显著抑制抗肿瘤免疫反应。淋巴瘤源性外泌体通过多条途径促进肿瘤发生和进展。BART miRNA具有免疫抑制功能,诱导巨噬细胞向M2型极化。外泌体介导了BART miRNA从Epstein-Barr病毒(Epstein-Barr virus,EBV)感染细胞向未感染细胞转移,诱导EBV相关淋巴瘤的发生^[31]。大量研究证实淋巴瘤源性外泌体可促进基质细胞增殖、侵袭和血管新生,诱导T淋巴细胞内PD-1、CTLA-4和TGF- β 等免疫抑制性蛋白的上调,抑制抗肿瘤免疫反应^[32]。外泌体与淋巴瘤耐药也具有很大的相关性,研究发现外泌体内Bcl-6 mRNA水平与利妥昔单抗抗体的耐药性呈正相关,外泌体介导了这种耐药基因在淋巴瘤细胞之间的交流^[33,34]。

早期诊断、精准治疗是淋巴瘤目前研究的主要方向。目前国内外已经研发了数个基于外泌体的疾病诊断试剂盒,Exosome Diagnostics公司推出全球首个基于外泌体的液态活检产品,可准确检测非小细胞肺癌的EML4-ALK突变^[35,36]。许多研究表明淋巴瘤源性外泌体表面可表达有肿瘤细胞的特异性表面标志,包括c-Myc、Mcl-1和Bcl-2等,同时某些标志还能在一定程度上区分淋巴瘤的类型,比如B细胞淋巴瘤源性外泌体表达CD19和CD20,T细胞淋巴瘤表达CD3等^[37,38]。相对于实体肿瘤,淋巴瘤在发病初期即随血液循环到达全身各个部位,很难进行有效手术切除,同时这也意味着在外周血中淋巴瘤源性外泌体浓度更高,在发病早期即可检测到。因此外泌体在淋巴瘤早期诊断、进展和复发监测中具有广阔的应用前景^[4]。Feng等^[37]对弥漫性大B细胞淋巴瘤进行基因测序,筛选出了显著上调的miR共37种,下调的miR共17种,并进一步分析了其与治疗效果和临床病理的相关性,发现血浆外泌体miR-125b-5p和miR-99a-5p可有效预测淋巴瘤的化疗效果,且与患者的生存时间紧密相关。Provencio M等^[39]分析了近100例淋巴瘤患者的临床标本,发现c-Myc、Bcl-6、Bcl-x1、AKT和PTEN可作为生物标志物监测淋巴瘤的治疗效果,预测患者的预后。

鉴于外泌体在淋巴瘤中的重要作用,细胞实验和动物实验结果均显示抑制外泌体分泌可显著抑制淋巴瘤生长、转移和耐药。然而这种方法基本不可能应用于临床,因为外泌体在维持人体稳态平衡和各系统功能中发挥着至关重要的作用。外泌

体属于双重生物膜构成的纳米囊泡样结构,科学家很容易对其进行材料学修饰^[40]。通过改造供体细胞或者在外泌体表面组装靶向分子,可增强外泌体治疗的靶向性。淋巴瘤表面特异性表达 c-Myc、Mc l-1 和 Bcl-2 等分子,通过聚乙烯亚胺(polyethylenimine, PEI)等修饰方法将相应抗体结合到外泌体表面,体内和体外结果均显示修饰后的外泌体特异性结合淋巴瘤细胞的能力显著升高^[41,42]。通过电转染或者化学转染的方式将 DNA、RNA 或者小分子递送到外泌体内,增强外泌体治疗的作用效能^[43]。研究者将阿霉素包裹到外泌体内,克服了化疗药物无法通过血脑屏障问题,可靶向杀伤脑组织内淋巴瘤细胞,药物作用时间延长,且副作用也显著降低,这为中枢神经系统肿瘤治疗提供了新的手段^[44]。

5 结语

淋巴瘤是最常见的血液系统恶性肿瘤,早期诊断和精准治疗具有十分重要的临床意义。外泌体是近几年研究的热点,大量研究证实外泌体与恶性肿瘤的发生、进展和耐药性紧密相关。外泌体上特异性标志和 miRNA 可用于淋巴瘤的早期诊断、复发监测和治疗效果预测。材料学修饰可显著增强外泌体治疗的靶向性和治疗效能。当然这些方法的有效性还需要大规模的临床数据评估。筛选高度特异性生物标志物、构建基于外泌体的靶向治疗体系、外泌体与疾病相关性研究是未来外泌体的主要发展方向。相信随着纳米材料、液体活检和基因技术的发展,在不久的将来外泌体产品将进入临床应用于淋巴瘤诊断和治疗。

参考文献(References)

- [1] Kawano N, Nagahiro Y, Yoshida S, et al. Clinical features and treatment outcomes of opportunistic infections among human T-lymphotrophic virus type 1 (HTLV-1) carriers and patients with adult T-cell leukemia-lymphoma (ATL) at a single institution from 2006 to 2016[J]. J Clin Exp Hematop, 2019, 59(4): 156-67
- [2] Meng J, Pan H, Li X, et al. Diagnosis and Treatment of Synchronous Lymphoma and Digestive System Carcinoma: Report of Four Cases and Literature Review[J]. Front Oncol, 2019, 9: 1367
- [3] Thodou E, Befani M, Triantafyllidis G, et al. Hodgkin Lymphoma Mimicking Inflammatory Breast Carcinoma: A Rare Case with Diagnostic Challenge and Novel Treatment [J]. Case Rep Hematol, 2019, 2019: 9256807
- [4] Zhang W, Zhu J, He X, et al. Exosome complex genes mediate RNA degradation and predict survival in mantle cell lymphoma[J]. Oncol Lett, 2019, 18(5): 5119-28
- [5] Nishio M, Teranishi Y, Morioka K, et al. Real-time assay for exosome membrane fusion with an artificial lipid membrane based on enhancement of gramicidin A channel conductance [J]. Biosens Bioelectron, 2020, 150: 111918
- [6] Lu Y, Duan Y, Xu Q, et al. Circulating exosome-derived bona fide long non-coding RNAs predicting the occurrence and metastasis of hepatocellular carcinoma[J]. J Cell Mol Med, 2020, 24(2): 1311-8
- [7] Zhao L, Gu C, Gan Y, et al. Exosome-mediated siRNA delivery to suppress postoperative breast cancer metastasis[J]. J Control Release, 2019, 318: 1-15
- [8] Wu Y, Liu W, Chen J, et al. Nuclear Exosome Targeting Complex Core Factor Zcchc8 Regulates the Degradation of LINE1 RNA in Early Embryos and Embryonic Stem Cells [J]. Cell Rep, 2019, 29(8): 2461-7
- [9] Mathews P M, Levy E. Exosome Production Is Key to Neuronal Endosomal Pathway Integrity in Neurodegenerative Diseases[J]. Front Neurosci, 2019, 13: 1347
- [10] Wang J, Lv B, Su Y, et al. Exosome-Mediated Transfer of lncRNA HOTTIP Promotes Cisplatin Resistance in Gastric Cancer Cells by Regulating HMGA1/miR-218 Axis [J]. Onco Targets Ther, 2019, 12: 11325-38
- [11] Anticoli S, Manfredi F, Choiozzini C, et al. An Exosome- Based Vaccine Platform Imparts Cytotoxic T Lymphocyte Immunity Against Viral Antigens[J]. Biotechnol J, 2018, 13(4): e1700443
- [12] Borrelli C, Ricci B, Vulpis E, et al. Drug-Induced Senescent Multiple Myeloma Cells Elicit NK Cell Proliferation by Direct or Exosome-Mediated IL15 Trans-Presentation [J]. Cancer Immunol Res, 2018, 6(7): 860-9
- [13] Rayamajhi S, Nguyen T D T, Marasini R, et al. Macrophage- derived exosome-mimetic hybrid vesicles for tumor targeted drug delivery[J]. Acta Biomater, 2019, 94: 482-94
- [14] Khalyfa A, Kheirandish-gozal L, Gozal D. Exosome and Macrophage Crosstalk in Sleep-Disordered Breathing-Induced Metabolic Dysfunction[J]. Int J Mol Sci, 2018, 19(11): e3383
- [15] Zhang B, Yeo R W Y, Lai R C, et al. Mesenchymal stromal cell exosome-enhanced regulatory T-cell production through an antigen-presenting cell-mediated pathway [J]. Cytotherapy, 2018, 20(5): 687-96
- [16] Ren R, Tan X H, Zhao J H, et al. Bone marrow mesenchymal stem cell-derived exosome uptake and retrograde transport can occur at peripheral nerve endings [J]. Artif Cells Nanomed Biotechnol, 2019, 47(1): 2918-29
- [17] 丁静雅, 秦露丹, 徐勇, 等. Graves 病患者外周血中 CD4⁺CD25⁺Treg 细胞上的 PD-1/PD-L1 的表达及意义[J]. 现代生物医学进展, 2019, 19(4): 718-723
- [18] Yue Y, Wang C, Benedict C, et al. Interleukin-10 Deficiency Alters Endothelial Progenitor Cell-Derived Exosome Reparative Effect on Myocardial Repair via Integrin-Linked Kinase Enrichment [J]. Circ Res, 2020, 126(3): 315-29
- [19] Stamatikos A, Knight E, Vojtech L, et al. Exosome-Mediated Transfer of Anti-miR-33a-5p from Transduced Endothelial Cells Enhances Macrophage and Vascular Smooth Muscle Cell Cholesterol Efflux[J]. Hum Gene Ther, 2020
- [20] Xu Z H, Miao Z W, Jiang Q Z, et al. Brain microvascular endothelial cell exosome-mediated S100A16 up-regulation confers small-cell lung cancer cell survival in brain[J]. FASEB J, 2019, 33(2): 1742-57
- [21] Yang T, Fogarty B, Laforge B, et al. Delivery of Small Interfering RNA to Inhibit Vascular Endothelial Growth Factor in Zebrafish Using Natural Brain Endothelia Cell-Secreted Exosome Nanovesicles for the Treatment of Brain Cancer[J]. AAPS J, 2017, 19(2): 475-86
- [22] Zhang W, Zhou Q, Wei Y, et al. The exosome-mediated PI3k/Akt/mTOR signaling pathway in cervical cancer [J]. Int J Clin Exp Pathol, 2019, 12(7): 2474-84
- [23] Soeda N, Iinuma H, Suzuki Y, et al. Plasma exosome-encapsulated

- microRNA-21 and microRNA-92a are promising biomarkers for the prediction of peritoneal recurrence in patients with gastric cancer [J]. *Oncol Lett*, 2019, 18(5): 4467-80
- [24] Huang J, Shen M, YAN M, et al. Exosome-mediated transfer of miR-1290 promotes cell proliferation and invasion in gastric cancer via NKD1 [J]. *Acta Biochim Biophys Sin (Shanghai)*, 2019, 51(9): 900-7
- [25] Zhao K, Wang Z, Li X, et al. Exosome-mediated transfer of CLIC1 contributes to the vincristine-resistance in gastric cancer [J]. *Mol Cell Biochem*, 2019, 462(1-2): 97-105
- [26] Xiao F, Xiao S, Xue M. miR-139 Controls Viability Of Ovarian Cancer Cells Through Apoptosis Induction And Exosome Shedding Inhibition By Targeting ATP7A [J]. *Onco Targets Ther*, 2019, 12: 10727-37
- [27] Zhang H D, Jiang L H, Hou J C, et al. Exosome: a novel mediator in drug resistance of cancer cells[J]. *Epigenomics*, 2018, 10(11): 1499-509
- [28] Zhang C, Ji Q, Yang Y, et al. Exosome: Function and Role in Cancer Metastasis and Drug Resistance [J]. *Technol Cancer Res Treat*, 2018, 17: 1533033818763450
- [29] Cypryk W, Nyman T A, Matikainen S. From Inflammasome to Exosome-Does Extracellular Vesicle Secretion Constitute an Inflammasome Dependent Immune Response? [J]. *Front Immunol*, 2018, 9: 2188
- [30] Que R S, Lin C, Ding G P, et al. Increasing the immune activity of exosomes: the effect of miRNA-depleted exosome proteins on activating dendritic cell/cytokine-induced killer cells against pancreatic cancer[J]. *J Zhejiang Univ Sci B*, 2016, 17(5): 352-60
- [31] Higuchi H, Yamakawa N, Imadome K I, et al. Role of exosomes as a proinflammatory mediator in the development of EBV-associated lymphoma[J]. *Blood*, 2018, 131(23): 2552-67
- [32] Tosetti F, Vene R, CAMODECA C, et al. Specific ADAM10 inhibitors localize in exosome-like vesicles released by Hodgkin lymphoma and stromal cells and prevent sheddase activity carried to bystander cells[J]. *Oncoimmunology*, 2018, 7(5): e1421889
- [33] Koch R, Demant M, Aung T, et al. Populational equilibrium through exosome-mediated Wnt signaling in tumor progression of diffuse large B-cell lymphoma[J]. *Blood*, 2014, 123(14): 2189-98
- [34] Xiao XB, Gu Y, Sun DL, et al. Effect of rituximab combined with chemotherapy on the expression of serum exosome miR-451a in patients with diffuse large b-cell lymphoma [J]. *Eur Rev Med Pharmacol Sci*, 2019, 23(4): 1620-5
- [35] Okuma Y, Morikawa K, Tanaka H, et al. Prospective exosome-focused translational research for afatinib study of non-small cell lung cancer patients expressing EGFR (EXTRA study)[J]. *Thorac Cancer*, 2019, 10(2): 395-400
- [36] Castellanos-Rizaldos E, Zhang X, Tadigotla VR, et al. Exosome-based detection of activating and resistance EGFR mutations from plasma of non-small cell lung cancer patients[J]. *Oncotarget*, 2019, 10(30): 2911-20
- [37] Feng Y, Zhong M, Zeng S, et al. Exosome-derived miRNAs as predictive biomarkers for diffuse large B-cell lymphoma chemotherapy resistance[J]. *Epigenomics*, 2019, 11(1): 35-51
- [38] Nanbo A, Katano H, Kataoka M, et al. Infection of Epstein(-)Barr Virus in Type III Latency Modulates Biogenesis of Exosomes and the Expression Profile of Exosomal miRNAs in the Burkitt Lymphoma Mutu Cell Lines[J]. *Cancers (Basel)*, 2018, 10(7): e237
- [39] Provencio M, Rodriguez M, Cantos B, et al. mRNA in exosomes as a liquid biopsy in non-Hodgkin Lymphoma: a multicentric study by the Spanish Lymphoma Oncology Group [J]. *Oncotarget*, 2017, 8(31): 50949-57
- [40] Su MJ, Aldawsari H, Amiji M. Pancreatic Cancer Cell Exosome-Mediated Macrophage Reprogramming and the Role of MicroRNAs 155 and 125b2 Transfection using Nanoparticle Delivery Systems[J]. *Sci Rep*, 2016, 6: 30110
- [41] Alhasan AH, Patel PC, Choi CH, et al. Exosome encased spherical nucleic acid gold nanoparticle conjugates as potent microRNA regulation agents[J]. *Small*, 2014, 10(1): 186-92
- [42] Lee YS, Jung WY, Heo H, et al. Exosome-Mediated Ultra- Effective Direct Conversion of Human Fibroblasts into Neural Progenitor-like Cells[J]. *ACS Nano*, 2018, 12(3): 2531-8
- [43] Cao Y, Wu T, Zhang K, et al. Engineered Exosome-Mediated Near-Infrared-II Region V2C Quantum Dot Delivery for Nucleus-Target Low-Temperature Photothermal Therapy[J]. *ACS Nano*, 2019, 13(2): 1499-510
- [44] Li J, Tian T, Zhou X. The role of exosomal shuttle RNA (esRNA) in lymphoma[J]. *Crit Rev Oncol Hematol*, 2019, 137: 27-34

(上接第 2752 页)

- [25] Slowinski T, Morgera S, Joannidis M, et al. Safety and efficacy of regional citrate anticoagulation in continuous venovenous hemodialysis in the presence of liver failure: the Liver Citrate Anticoagulation Threshold (L-CAT) observational study[J]. *Crit Care*, 2015, 19: 349
- [26] 马元吉, 陈芳, 许艳, 等. 局部枸橼酸抗凝在血浆吸附联合血浆置换治疗高危出血倾向肝衰竭患者中的应用[J]. *世界华人消化杂志*, 2018, 26(3): 165-173
- [27] Liu N, Kan J. Effect of local sodium citrate anticoagulation on blood purification in patients with high-risk bleeding in ICU [J]. *Modern Medicine Journal of China*, 2019, 41(12): 1334-1339
- [28] Jiang W, Lin G, Shi H, et al. Clinical Application of Local Anticoagulation of Sodium Citrate in Blood Purification of Intensive Care Unit[J]. *Clinical Medicine & Engineering*, 2019, 26(7): 899-900
- [29] Huang Y L, Chen Y H, Yuan L, et al. Application of local anticoagulation with citrate in hemodialysis of patients with high risk of bleeding tendency[J]. *Journal of Clinical & Experimental Medicine*, 2019, 20(2): 94
- [30] 李春艳, 焦军东. 关于局部枸橼酸抗凝技术在合并肝损伤的血液净化患者中应用的研究进展 [J]. *中国中西医结合肾病杂志*, 2018, 19(11): 91-92